



MICROSHIELD® Tincture 2E

Composition

Chlorhexidine Gluconate Solution IP 10 % v/v (equivalent to Chlorhexidine Gluconate 2 % w/v)
Ethanol IP 80% v/v

Applications

- For Pre-Surgical and Post-Surgical Skin Antisepsis

Direction for use

- Use undiluted
- Swab Skin to visibly wet area to be treated
- Leave for 2 minutes
- Allow to dry

Instructions for use

Solutions are not for ophthalmic, bladder and peritoneal usage. Microshield® Tincture 2E must not come in contact with brain, meninges, middle ear or syringes and needles to be used for spinal injections.

Do not mix with detergents or other chemicals. Keep out of eyes and ears. Keep out of reach of children. Contraindicated in case of known hypersensitivity to Chlorhexidine.

Product Data

Contains Chlorhexidine Gluconate 2.0% w/v in 80% v/v Ethanol.IP.

Special advice

Always read the label and use only as directed.

- For external use only. Do not swallow.
- Flammable. Keep away from fire or flame.
- Avoid contact with eyes. In case of eye contact, immediately flush with water
- Discontinue use if skin irritation or redness develops.
- Do not mix with detergents and other chemicals.
- Do not use after the expiry date.
- Contraindicated in case of known hypersensitivity to Chlorhexidine.

Information for ordering

Item	Delivery form	Item no.
MICROSHIELD Tincture 2E100ml	100/Carton	70002449
MICROSHIELD Tincture 2E 500ml	20/Carton	70002450